

PAUTES DE CORRECCIÓ
SÈRIE 4FÍSICA
CURS 2005-06

P1. a) $W = M g (H-h)$ [0,7] $\rightarrow W = 300 \cdot 10 \cdot (15-5) = 3 \cdot 10^4 \text{ J}$ [0,3]


b) $W_{nc} = \Delta E = \Delta(u + E_c)$ [0,7]

$$W_{nc} = -30.000 + \frac{1}{2} 300 (10^2 - 0^2) = -15.000 \text{ J}$$

$$Q = -1,5 \cdot 10^4 \text{ J}$$
 [0,3]

c) $N - M g = M \frac{v^2}{R}$ [0,7]

$$N = M \left(g + \frac{v^2}{R} \right) \rightarrow N = 8 \cdot 10^3 \text{ N}$$
 [0,3]



Q1. a) $\theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2$ [0,2] $\rightarrow \alpha = \frac{2\theta}{t^2} = \frac{\pi}{40} \text{ rad/s}^2$ [0,1]

$$\omega = \omega_0 + \alpha t$$
 [0,1] $\rightarrow \omega = \frac{\pi}{40} \cdot 15 \text{ rad/s}$

$$a_n = \omega^2 r$$
 [0,1] $\rightarrow a_n = \left(\frac{\pi}{40} \cdot 15 \right)^2 \cdot 0,15 = 0,21 \text{ m/s}^2$ [0,1]

b) $a_t = \alpha \cdot r$ [0,3] $\rightarrow a_t = \frac{\pi}{40} \cdot 0,15 = 1,2 \cdot 10^{-2} \text{ m/s}^2$

[0,1]

Q2. $g_L/5 = G \frac{M_L}{(R_L+h)^2}$ [0,2] $\left. \begin{array}{l} \\ \\ \end{array} \right\} 5 = \left(\frac{R_L+h}{R_L} \right)^2 \rightarrow h = R_L (\sqrt{5}-1)$ [0,4]

$$g_L = G \frac{M_L}{R_L^2}$$
 [0,2]

$$h = 2,15 \cdot 10^6 \text{ m}$$
 [0,2]

OPCIÓ A

P2. a) $A = 0,01 \text{ m}$ [0,2]

$$k = 2\pi/\lambda = 2\pi\nu/v \rightarrow k = 2\pi \cdot \frac{4}{3} \text{ rad/m}$$
 [0,2]

$$\omega = 2\pi\nu \rightarrow \omega = 2\pi \cdot 440 \text{ rad/s}$$
 [0,2]

$$\rightarrow \psi = 0,01 \cdot \cos 2\pi \left(\frac{4}{3} x - 440 t \right)$$
 [0,4]

la solució en sin també és vàlida


b) $\phi = k \cdot \Delta x$ [0,6] $\rightarrow \phi = 5\pi \text{ rad}$ [0,2]

Defasatge real: $\pi \text{ rad}$. [0,2]

SÈRIE 4 (CONT.)

CURS 2005-06

$$c) v_{\max} = +Aw \quad [0,6] \rightarrow v_{\max} = \boxed{27,65 \text{ m/s}} \quad [0,4]$$

Totes les forces són d'atracció! [0,4]La resultant té la direcció de la diagonal del quadrat i sentit cap al centre. [0,6]


$$Q4. \quad \left. \begin{array}{l} E = h\nu \quad [0,2] \\ \lambda = c/\nu \quad [0,2] \end{array} \right\} E = \frac{hc}{\lambda} \quad [0,1] \rightarrow E = \boxed{3,3 \cdot 10^{-19} \text{ J}} \quad [0,1]$$

$$p = \frac{h}{\lambda} \quad [0,3] \rightarrow p = \boxed{1,1 \cdot 10^{-27} \frac{\text{kg} \cdot \text{m}}{\text{s}}} \quad [0,2]$$

Opció B

P2. a) $V = k \left(\frac{q}{a/2} + \frac{q}{a/2} + \frac{2q}{\sqrt{a^2 - \frac{a^2}{4}}} \right) \quad [0,7]$

$$V = k \frac{q}{a} \left(2 + 2 + \frac{4}{\sqrt{3}} \right) = \boxed{3,28 \cdot 10^6 \text{ V}} \quad [0,3]$$



$$b) \vec{E} = k \frac{q}{\left(\frac{a}{2}\right)^2} (1,0) + k \frac{q}{\left(\frac{a}{2}\right)^2} (-1,0) + k \frac{2q}{a^2 - \frac{a^2}{4}} (0,-1) \quad [0,3]$$

$$\rightarrow \vec{E} = \boxed{8 \cdot 10^5 (0,-1) \text{ N/C}} \quad [0,4]$$

$$c) \left. \begin{array}{l} W = 2q \cdot (V_f - V_i) \quad [0,3] \\ V_i = k \left(\frac{q}{a} + \frac{q}{a} \right) \quad [0,2] \\ V_f = k \left(\frac{q}{a/2} + \frac{q}{a/2} \right) \quad [0,2] \end{array} \right\} W = 2q \cdot k \frac{2q}{a} = \boxed{2,1 \text{ J}} \quad [0,3]$$

Q3. 1.a, 2.b

Q4. 1.c, 2.a

Correcta: 0,5En blanc: 0Incorrecta: -0,25

} El total de Q3+Q4
entre 0 i 2 punts
(no puntuacions
negatives)